

Arsenic-containing hydrocarbons are natural constituents of sashimi tuna

Mojtaba S. Taleshi¹, John S. Edmonds¹, Walter Goessler¹, Maria José Ruiz-Chancho¹, Georg Raber¹, Kenneth B. Jensen², Kevin A. Francesconi¹

¹ Institute of Chemistry–Analytical Chemistry Karl-Franzens University Graz,
Universitaetsplatz 1, 8010 Graz (Austria)

² Institute for Physics and Chemistry, University of Southern Denmark, 55 Campusvej, 5230
Odense, Denmark

Arsenic occurs naturally in many types of seafood as water-soluble and fat-soluble organoarsenic compounds. Although water-soluble compounds have been well characterised, the fat-soluble compounds, so-called arsenolipids, have until recently remained unknown. We report that sashimi grade tuna fish, with a total arsenic content of 5.9 µg As/g dry mass, contains approximately equal quantities of water-soluble and fat-soluble arsenic. The water-soluble arsenic was predominantly arsenobetaine. Two fat-soluble compounds were isolated and characterised. The first was identified as 1-dimethylarsinylpentadecane [(CH₃)₂As(O)(CH₂)₁₄CH₃] by comparison of HPLC/mass spectrometric data and accurate mass data with those of an authenticated synthesised standard. The second arsenolipid was postulated as 1-dimethylarsinyl *all-cis*-4,7,10,13,16,19-docosahexane from mass spectrometric data and analogy with non arsenic-containing lipids found in fish. These two arsenolipids constituted about 50% of the total fat-soluble arsenic; the other 50% consisted of less polar arsenolipids of currently unknown structure. This is the first identification of an arsenolipid in commonly consumed seafood.